# **Cervix Uteri Cancer Staging**

7th EDITION

## **Definitions**

#### **Primary Tumor (T)**

Primar	y Tum	or (T)		
TNM CATEGORIES	FIGO STAGES			
TX		Primary tumor cannot be assessed		
T0		No evidence of primary tumor		
Tis*		Carcinoma in situ (preinvasive carcinoma)		
T1	I	Cervical carcinoma confined to uterus (extension to corpus should be disregarded)		
T1a**	IA	Invasive carcinoma diagnosed only by microscopy. Stromal invasion with a maximum depth of 5.0 mm measured from the base of the epithelium and a horizontal spread of 7.0 mm or less. Vascular space involvement, venous or lymphatic, does not affect classification		
T1a1	IA1	Measured stromal invasion 3.0 mm or less in depth and 7.0 mm or less in horizontal spread		
T1a2	IA2	Measured stromal invasion more than 3.0 mm and not more than 5.0 mm with a horizontal spread 7.0 mm or less		
T1b	IB	Clinically visible lesion confined to the cervix or microscopic lesion greater than T1a/IA2		
T1b1	IB1	Clinically visible lesion 4.0 cm or less in greatest dimension		
T1b2	IB2	Clinically visible lesion more than 4.0 cm in greatest dimension		
<b>T2</b>	II	Cervical carcinoma invades beyond uterus but not to pelvic wall or to lower third of vagina		
T2a	IIA	Tumor without parametrial invasion		
T2a1	IIA1	Clinically visible lesion 4.0 cm or less in greatest dimension		
T2a2	IIA2	Clinically visible lesion more than 4.0 cm in greatest dimension		
T2b	IIB	Tumor with parametrial invasion		
ТЗ	III	Tumor extends to pelvic wall and/or involves lower third of vagina, and/or causes hydronephrosis or nonfunctioning kidney		
T3a	IIIA	Tumor involves lower third of vagina, no extension to pelvic wall		
T3b	IIIB	Tumor extends to pelvic wall and/or causes hydronephrosis or nonfunctioning kidney		
<b>T4</b>	Tumor invades mucosa of bladder or rectum, and/or extends beyond true pelvis (bullous edema is not sufficient to classify a tumor as T4)			


#### **Regional Lymph Nodes (N)**

TNM CATEGORIES	FIGO STAGES	
NX		Regional lymph nodes cannot be assessed
NO		No regional lymph node metastasis
N1	IIIB	Regional lymph node metastasis

### **Distant Metastasis (M)**


TNM ATEGORIES	FIGO STAGES	
MO	STAGES	No distant metastasis
M1	IVB	Distant metastasis
		(including peritoneal
		spread, involvement
		of supraclavicular,
		mediastinal, or paraaorti
		lymph nodes, lung,
		liver, or bone)


ANATOMIC STAGE/PROGNOSTIC GROUPS (FIGO 2008)						
Stage 0*	Tis	N0	M0			
Stage I	T1	N0	M0			
Stage IA	T1a	N0	M0			
Stage IA1	T1a1	N0	M0			
Stage IA2	T1a2	N0	M0			
Stage IB	T1b	N0	M0			
Stage IB1	T1b1	N0	M0			
Stage IB2	T1b2	N0	M0			
Stage II	T2	N0	M0			
Stage IIA	T2a	N0	M0			
Stage IIA1	T2a1	N0	M0			
Stage IIA2	T2a2	N0	M0			
Stage IIB	T2b	N0	M0			
Stage III	T3	N0	M0			
Stage IIIA	T3a	N0	M0			
Stage IIIB	T3b	Any N	M0			
	T1-3	N1	M0			
Stage IVA	T4	Any N	M0			
Stage IVB	Any T	Any N	M1			



### **Notes**

- \*FIGO no longer includes Stage 0 (Tis).
- \*\*\* All macroscopically visible lesions—even with superficial invasion—are T1b/IB.







Financial support for AJCC 7th Edition Staging Posters provided by the American Cancer Society

